

NINKOKO

Festuca arundinacea

- Late heading with very high total dry matter yield
- Excellent resistance to rust and other diseases
- Superior drought tolerance
- 51 days spring window for optimal use



High dry matter production

French official trials show high total dry matter production with a high first cut in the spring.

Spring window

With 51 days between 30 cm grass in the spring, and the beginning of ear emergence, this late heading variety gives flexibility in the use without losing quality.

Disease tolerance and softness of leaves

With a score of 6.4 for softness of leaves, Ninkoko joins the group of soft leaved varieties.

The excellent rust tolerance, with a score of 7.5 adds quality and palatability to the autumn and summer production.

Technical Specifications


- Tall fescue
- Top yielding with high disease tolerance
- Listed/recommended in EU CZ FR
- Lateness: Late

Specification and ratings

Dry matter yield

Source : Geves 2015 (France)

Dry matter yield in t/ha



Characteristics

Source: Geves 2015 (France)

Variety	Ninkoko	Average catalogue
Heading date	6 May	6 May
30 cm grass height in spring	19 March	19 March
Spring window in days	51,0	50,3
Softness of leaves 1-9, 9 = softer	6,4	6,5
Rust tolerance 1-9, 9 = tolerant	7,5	6,9

Ratings

Scale 1-9, where 9 = best or most pronounced

